

Included-row exchange principle for quantum minors

Zoran Škoda*

February 2, 2008

¹ Bialgebra $\mathcal{M}_q(n)$ ('the quantum linear semigroup') and its Hopf quotients $\mathcal{SL}_q(n)$, $\mathcal{GL}_q(n)$ and their multiparametric analogues have been studied in a number of references with diverse motivation. Identities among the quantum minors in those algebras were important in numerous calculations, often with representation-theoretic and geometric motivation. Though more complicated identities will still appear, the subject is considered matured in a sense: if we consider only homogeneous order two (counting minors involved) relations and impose certain natural ordering on the minors, in a sense all such "commutation" relations are known. The situation is more complicated with higher order relations and when the order of labels in identities is non-standard.

In geometrical study of the algebras above, quantum Gauss decomposition plays a major role. Unlike in the classical case, the shifts of the big Bruhat cell are not mutually isomorphic (as algebras). Depending on formalism, the permutation of rows or columns in the Gauss decomposition problem forces different relations among the generators of the solutions. However, some among the related identities surprisingly stay the same, although the commutation relations and hence the proofs of the relations change significantly after the ordering among the rows changes.

In this paper we prove a principle for generalizing certain identities, where a position of a row of exactly one of the entries involved in each monomial changes, without changing the shape of the identity, including the exact coefficients. We call this the **included-row exchange principle** for identities

*Theoretical Physics Division, Institute Rudjer Bošković, Bijenička cesta 54, P.O.Box 180, HR-10002 Zagreb, Croatia; zskoda@irb.hr

involving quantum minors because the position of the exchanged rows is included in all other minors appearing in the identity.

2 Relations in $\mathcal{M}_q(n)$. Let K, L be some linearly ordered sets of positive cardinality. The elements of K and L will be called labels. The generators of associative unital algebra $\mathcal{M}_q(K, L)$ over a base field $\mathbf{k}$ are t_l^k where $k \in K$, $l \in L$, $0 \neq q \in \mathbf{k}$ and the relations are

$$\begin{aligned} t_\gamma^\alpha t_\delta^\alpha &= q t_\delta^\alpha t_\gamma^\alpha, & \alpha = \beta, \gamma < \delta \text{ (same row)} \\ t_\gamma^\alpha t_\gamma^\beta &= q t_\gamma^\beta t_\gamma^\alpha, & \alpha < \beta, \gamma = \delta \text{ (same column)} \\ [t_\gamma^\alpha, t_\delta^\beta] &= (q - q^{-1}) t_\gamma^\beta t_\delta^\alpha (\theta(\delta > \gamma) - \theta(\alpha > \beta)), & \alpha \neq \beta \text{ and } \gamma \neq \delta \end{aligned}$$

where $\theta(\text{true}) = 1$, $\theta(\text{false}) = 0$, and $[,]$ stands for the ordinary commutator. In particular, the isomorphism class of algebra $\mathcal{M}_q(K, L)$ depends only on the cardinality of K and L , and we denote a representative of that class $\mathcal{M}_q(n)$ if $|K| = |L| = n$.

Clearly, for each pair of inclusions of ordered sets $K \hookrightarrow K', L \hookrightarrow L'$, there is a canonical inclusion of associative algebras $\mathcal{M}_q(K, L) \hookrightarrow \mathcal{M}_q(K', L')$. For convenience, in the same spirit, we will embed $\mathcal{M}_q(n)$ into some algebra $\mathcal{M}_q(\infty)$, which is defined the same way, except that the sets of labels K and L are countably infinite.

3 Definition. The quantum exterior algebra $\Lambda_q(n)$ is an associative unital algebra on n generators, $e_1, \dots, e_n$ and multiplication will be denoted by $\wedge$ modulo the relations

$$e_i \wedge e_j = -q^{-1} e_j \wedge e_i, \quad i < j,$$

and $e_i^2 = 0$ for all i .

Lemma. A vector space basis of $\Lambda_q(n)$ is given by the set of all sequences $e_{i_1} e_{i_2} \dots e_{i_m}$ where $0 \leq m \leq n$ and $i_1 < i_2 < \dots < i_m$. The proof is by diamond lemma.

Definition. The left canonical coaction δ_l and the right canonical coaction δ_r of $\mathcal{O}(M_q(n))$ on $\Lambda_q(n)$ are the unique homomorphisms of module algebras for which

$$\delta_r(e_i) = \sum_j e_j \otimes t_i^j, \quad \delta_l(e_i) = \sum_j t_j^i \otimes e_j$$

Then $\delta_r(e_1 \wedge \dots \wedge e_n) = e_1 \wedge \dots \wedge e_n \otimes c_r$ and analogously $\delta_l(e_1 \wedge \dots \wedge e_n) = c_l \otimes e_1 \wedge \dots \wedge e_n$ for certain elements $c_l, c_r \in \mathcal{O}(M_q(n))$.

Denote by $l(\sigma)$ the length (number of inversions) of the permutation $\sigma \in \Sigma(n)$.

Definition. Let $\sigma, \tau \in \Sigma(n)$. For ring $R = \mathcal{O}(M_q(n))$ define maps

$$d_\tau^{c,\sigma}, d_\tau^{r,\sigma}, d^c, d^r : M_n(R) \rightarrow R$$

by

$$\begin{aligned} d_\tau^{r,\sigma}(G) &= (-q)^{-l(\sigma)+l(\tau)} \sum_{\rho \in \Sigma(n)} (-q)^{l(\rho)} h_{\rho(1)}^1 \cdots h_{\rho(n)}^n \\ d_\tau^{c,\sigma}(G) &= (-q)^{l(\sigma)-l(\tau)} \sum_{\rho \in \Sigma(n)} (-q)^{l(\rho)} h_1^{\rho(1)} \cdots h_n^{\rho(n)} \end{aligned}$$

where $h_j^i = g^\sigma(i)_{\tau(j)}$. We skip σ and τ when they are equal to identity permutations.

Quantum determinant. Fact:

$$d_\tau^{r,\sigma}(T) = d_\tau^{c,\sigma}(T) = c_r = c_l$$

We denote that element D_q or $\det_q T \in \mathcal{O}(M_q(n))$ and call it quantum determinant. It is central and group-like.

For any pair of m -member subsets $K, L \subset \{1, \dots, n\}$ define the **quantum subdeterminants** (quantum minors)

$$D_L^K = \det_q T_L^K = \sum_{\sigma \in \Sigma(m)} (-q)^{l(\sigma)} t_{(c_L \circ \sigma)(1)}^{c_K(1)} \cdots t_{(c_L \circ \sigma)(m)}^{c_K(m)}.$$

where c_K, c_L are the counting functions, i.e. unique strictly monotone functions $\{1, \dots, m\} \rightarrow K$ and L .

Let $\phi : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ is any function. Then the matrix T^ϕ given by

$$(T^\phi)_j^i = T_j^{\phi(i)}$$

is called a permuted quantum matrix with repeated rows if ϕ is not injective.

Lemma. $d^r(T^\phi) = 0$ iff ϕ is not injective.

4 Laplace expansions. For any pair of m -member subsets $K, L \subset \{1, \dots, n\}$

$$\begin{aligned} \delta_L^K D_q &= \sum_J (-q)^{J-L} \det_q T_J^K \det_q T_{\hat{J}}^L \\ &= \sum_J (-q)^{J-L} \det_q T_K^J \det_q T_{\hat{L}}^J \\ &= \sum_J (-q)^{L-J} \det_q T_{\hat{J}}^L \det_q T_J^K \\ &= \sum_J (-q)^{L-J} \det_q T_{\hat{L}}^J \det_q T_K^J \end{aligned}$$

where J runs over m -member subsets of $\{1, \dots, n\}$ and the name of a subset in exponent denotes the sum of its elements. Also $\hat{}$ denotes a complement.

5 Suppose $D = D_L^K = \det_q T_L^K$ is some quantum minor, $|K| = |L|$. Let $k \in K, k' \notin K$. Then $D_L^{K(k \rightarrow k')} := \det_q T_L^{K \setminus \{k\} \cup \{k'\}}$. Similarly, one defines $D_{L(l \rightarrow l')}^K$ and iterations like $D_L^{K(k, k'' \rightarrow k', k''')} = D_L^{K(k \rightarrow k')(k'' \rightarrow k''')}$ etc.

6 We will consider the identities involving quantum minors D_L^K (including the 1×1 case, where $D_l^k = t_l^k$). Formally, we form a free algebra $\mathcal{F}$ on generators Δ_L^K where K, L are multilabels with $|K| = |L|$. Let $\pi : \mathcal{F} \rightarrow \mathcal{M}_q$ be the natural projection, where $\pi(\Delta_L^K) = D_L^K$. We also denote $f_j^i := \Delta_j^i$. An **identity among quantum minors** is an element $f \in \mathcal{F}$ such that $\pi(f) = 0$. The identity will be called **homogeneous** if for each r , the number of factors of D_L^K with $|K| = |L| = r$ is the same for each monomial in $\mathcal{F}$.

We would like to do certain replacements of labels in identities.

7 Let $\mathcal{F}_N$ be the bigger free algebra on the 'decorated' symbols $\Delta_L^K(i)$ where $i = 0, 1, 2, \dots$. Note the canonical 0-th inclusion $\iota_0 : \mathcal{F} \hookrightarrow \mathcal{F}_N$ $\Delta_L^K \mapsto \Delta_L^K(0)$ and the canonical projection $\pi_N : \mathcal{F}_N \rightarrow \mathcal{F}$ given by $\Delta_L^K(i) \rightarrow \Delta_L^K$ for all i . A **decorated identity** among quantum minors is an element $f \in \mathcal{F}_N$ so that $\pi(\pi_N(f)) = 0$.

For that purpose, we take a finite sequence $A = (A_1, \dots, A_r)$ where, for each $1 \leq i \leq r$, $A_i = (K_i, L_i \rightarrow K'_i, L'_i)$ is some nontrivial 'rule of replacement of labels', where possibly K -s or L -s (but not both) may be ommitted. Formally we may also put $A_i = 0$ for $i = 0$ and for $i > r$. There is a unique endomorphism ϕ_A of $\mathcal{F}_N$ which sends $D_J^I(i)$ to $D_{J(L_i \rightarrow L'_i)}^{I(K_i \rightarrow K'_i)}(i)$ whenever $A_i = (K, L \rightarrow K', L')$, $K_i \subset I$ and $L_i \subset J$ and fixes (=does not change) $D_J^I(i)$ for those triples (i, I, J) for which these conditions fail. If K -s are dropped than we do changes for all A decorated element $f \in \mathcal{F}_N$ is an **injective match** for A if each monomial in f has for each $0 < i \leq r = r_A$ exactly one factor $D_J^I(i)$ (and that one appearing exactly once in the monomial) such that $K_i \subset I$ and $L_i \subset J$.

For example, let $A = ((2, 1 \rightarrow 3, 1), (33, 23 \rightarrow 23, 23))$ is an injective match for $f = \Delta_{23}^{33}(2)f_3^3(0)f_1^2(1)f_3^3(0)$ and $\phi_A(f) = \Delta_{33}^{23}(2)f_3^3(0)f_1^3(1)f_3^3(0)$.

8 Quantum Muir's law of extensionality. [5] *Let a homogeneous identity $\mathcal{I}$ between quantum minors of a submatrix T_J^I of a matrix T be given. Let $K \cap I = \emptyset$, $J \cap L = \emptyset$ and $K = J$. If every quantum minor D_V^U of T_J^I in the identity $\mathcal{I}$ is replaced by the quantum minor $D_{V \cup J}^{U \cup I}$ of $T_{J \cup L}^{I \cup K}$, then we obtain a new identity $\mathcal{I}'$ called *extensional* to $\mathcal{I}$. There is also a nonhomogeneous version involving pivot, and we won't use it; in a nutshell we multiply some summands with 1 treated as a quantum minor with an empty set of row indices and an empty set of column indices to get a formally homogeneous*

identity.

9 Theorem. *Let f be a decorated identity among quantum minors, let $k \neq k'$ be row labels, and l_0 a column label such that*

1) (LINEARITY) *all factors in each monomial of f are of the form $\Delta_L^K(0)$ except one. That one is $f_l^k(1)$ with $l \neq l_0$ and appears exactly once in each monomial. The label l may depend on the monomial involved.*

2) (INCLUDED ROW CONDITION) *row labels k, k' and column l_0 appear in each 0-type factor $\Delta_L^K(0)$ of each monomial f_i in f .*

3) (HIERARCHY OF IDENTITIES) *If we erase row k' and column l_0 from each zero type factor $\Delta_L^K(0)$, we get a decorated identity $f^\vee$.*

Conclusion: $A = (A_1) = ((k \rightarrow k'))$ injectively matches f and $\phi_A(f)$ is a decorated identity among quantum minors (said to be obtained from f by the included row exchange).

Proof. We first apply Muir's law to $f^\vee$ by extending each minor in $f^\vee$ by k' -th row and l_0 -column. The obtained identity $\tilde{f} = (f^\vee)_{+l_0}^{+k'}$ differs from f and from $\phi_A(f)$ respectively, in the sense that the factor $t_l^k(1)$, or $t_l^{k'}(1)$ respectively, is replaced by $\Delta_{ll_0}^{kk'}$. Now we project to $\mathcal{M}_q(n)$ to check that we have an identity. As $D_{ll_0}^{kk'} = t_l^k t_{l_0}^{k'} - q t_l^{k'} t_{l_0}^k$, by linearity 1) we can write identity $\tilde{f}$ as a sum of the two terms, $\tilde{f}^k$ and $\tilde{f}^{k'}$ where $\tilde{f}^k$ is obtained from $f_l^k f_{l_0}^{k'}$ summand in $\Delta_{ll_0}^{kk'}$.

Claim 1: $\tilde{f}^k - f \cdot f_{l_0}^{k'}(0)$ is a (decorated) identity. This follows from the included row condition 2), as k' -th row and l_0 -th column are included in each other factor. Hence in $\mathcal{M}_q(n)$, $t_{l_0}^{k'}$ commutes with each D_L^K involved and hence $f_{l_0}^{k'}$ can be pushed all to the right without change in the coefficients on the expense of an expression which is an identity.

Claim 2: $\tilde{f}^{k'} - q\phi_A(f) \cdot f_{l_0}^k(0)$ is a (decorated) identity. Again, by the included row condition 2), $t_{l_0}^k$ can be also pushed to the right end.

Now recalling that f is an identity, we obtain that $f \cdot f_{l_0}^{k'}(0)$ is an identity and by claim 1 it follows that $\tilde{f}^k$ is also an identity. As $\tilde{f}$ is also the identity, then $\tilde{f}^{k'} = \tilde{f} - \tilde{f}^k$, and, by claim 2, $q\phi_A(f) \cdot f_{l_0}^k(0)$ are identities too. $\mathcal{M}_q(n)$ is well-known to be a domain (no zero divisors), hence $\phi_A(f)$ is the identity, as required. This finishes the proof.

Examples (and applications related to quantum Gauss decomposition) will be treated elsewhere. The author has also used this technique in his Ph.D. University of Wisconsin thesis (defended January 2002).

References

- [1] M. ARTIN, W. SCHELTER, J. TATE, *Quantum deformations of $GL(n)$* , Comm. Pure Appl. Math. Vol. XLIV (1991), pp. 879–895.
- [2] E. E. DEMIDOV, *Multiparameter quantum deformations of the group $GL(n)$* . (Russian) *Uspehi Mat. Nauk* 46 (1991), no. 4(280), pp. 147–148; translation in *Russian Math. Surveys* 46 (1991), no. 4, pp. 169–171.
- [3] I.M. GEL’FAND, V.S. RETAKH, *Determinants of matrices over non-commutative rings*, Funct. Anal. Appl. vol 25 (1991), no. 2, pp. 91–102. transl. 21 (1991), pp. 51–58.
- [4] I. M. GEL’FAND, V. S. RETAKH, *Quasideterminants I*, Selecta Mathematica, New Series 3 (1997) no.4, pp. 517–546; q-alg/9705026.
- [5] D. KROB, B. LECLERC, *Minor identities for quasi-determinants and quantum determinants*, Comm.Math.Phys. 169 (1995), pp. 1–23.
- [6] YU. I. MANIN, *Quantum groups and non-commutative geometry*, CRM, Montreal 1988.
- [7] YU. I. MANIN, *Multiparametric quantum deformation of the general linear supergroup*, Comm. Math. Phys. 123 (1989), pp. 163–175.
- [8] B. PARSHALL, J. WANG, *Quantum linear groups*, Memoirs AMS 439, 1991.
- [9] Z. ŠKODA, *Localizations for construction of quantum coset spaces*, in “Noncommutative geometry and quantum groups”, P. M. Hajac, W. Pusz eds. Banach Center Publications vol.61, pp. 265–298, Warszawa 2003; arXiv:math.QA/0301090.